

Dengue Fever Associated Foveolitis

Sameeksha Agrawal, Ankit Agrawal, Krishna K. Agrawal, Vijay K. Agrawal

Department of Ophthalmology, Krishna Eye Foundation, Eye Clinic, Prayagraj, Uttar Pradesh, India.

Abstract

Purpose: To report a case of foveolitis associated with dengue fever.

Methods: Retrospective, observational case report.

Case description: A 12-year-old male child, presented to the retina department with complaints of sudden blurring of vision in left eye since 1 week following dengue fever. His presenting visual acuity in left eye was 20/60, N8. Fundus showed well-circumscribed, pale yellow-orange lesion at the foveal center in LE. Retinal imaging findings were consistent with the diagnosis of foveolitis.

Results: Foveolitis and macular swelling were found to resolve over time on serial OCT, with visual acuity improving to 20/20, N6 with a resolution of scotoma.

Conclusion: Infection with dengue virus might be associated with foveolitis. We suggest that serial OCT be performed to monitor maculopathy.

Keywords: Dengue fever, Maculopathy, Disruption of outer retina, Foveolitis, Y split in outer plexiform layer.

INTRODUCTION

Dengue fever is the most common mosquito-borne viral disease in humans that is transmitted through the bite of an infected female *Aedes aegypti/albopictus* mosquitoes and is commonly found in the tropics.¹

With the recent outbreak of dengue, India recorded 110473 dengue cases in 2022, according to data shared by the National Center for vector-borne Diseases Control.²

Dengue fever is a multisystemic disease with known complications. It may cause maculopathy with a wide spectrum of manifestations. Ocular findings were mostly seen in the posterior pole of the fundus, manifesting as retinal hemorrhages, macular edema, foveolitis, vasculitis, and optic neuropathy.^{3,4}

In this case report, we report a case of foveolitis following dengue fever which showed spontaneous recovery through the follow-up visits

Case Description

A 12-year-old male child presented to the retina department with complaints of sudden blurring of vision in left eye since one week following dengue fever. His presenting visual acuity in right eye was 20/20 and left eye was 20/60, N8. RE

fundus looked grossly normal at presentation (Figure 1). OCT line scan through fovea showed “Y split” in outer plexiform layer (Figure 2A). Although at 1 week followup. RE started showing parafoveal photoreceptor layer changes suggestive of photoreceptoritis. There was focal discontinuity in the ellipsoid zone and the external limiting membrane just above the intact retinal pigment epithelium (Figure 2B).

At presentation, LE fundus showed a well-circumscribed, pale yellow-orange lesion at the foveal center (Figure 1). The corresponding area of the focal outer neurosensory retina–RPE thickening at the foveal center was identified on OCT, suggestive of foveolitis. A hyperreflective conical lesion was noted arising beneath the ELM. Associated “Y-shaped” split can be seen in outer plexiform layer (Figure 3A). Amsler’s grid charting revealed a central scotoma in the LE. At day 7 post-

Address for correspondence: Sameeksha Agrawal

Department of Ophthalmology, Krishna Eye Foundation, Eye Clinic, Prayagraj, India.

E-mail: sameeksha.agrawal92@gmail.com

© UPIO, 2023 Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <https://creativecommons.org/licenses/by-nc-sa/4.0/>.



UP JOURNAL OF OPHTHALMOLOGY

An Official Journal of Uttar Pradesh State Ophthalmological Society,
UPSOS (Northern Ophthalmological Society, NOS)

p-ISSN: 2319-2062

DOI: 10.56692/upjo.2023110105

How to cite this article: Agrawal S, Agrawal A, Agrawal KK, Agrawal VK. Dengue Fever Associated Foveolitis. UP Journal of Ophthalmology. 2023;11(1): 23-25.

Received: 02-12-2022, **Accepted:** 16-01-2023, **Published:** 31-03-2023




Figure 1: Fundus picture of RE was grossly Normal and LE shows well-circumscribed, pale yellow-orange lesion at the foveal center




Figure 2: (A) At presentation – OCT line scan through fovea shows “Y split” in outer plexiform layer, (B) Comparative scan shows changes at photoreceptor layer at 1 week (Photoreceptoritis). There is focal discontinuity in the ellipsoid zone and the external limiting membrane just above the intact retinal pigment epithelium, (C) Shows 1 month follow-up scan, photoreceptoritis almost resolved, (D) Complete resolution seen at 2 month follow-up.

presentation, the imaging findings showed improvement in LE with resolution of conical foveal lesion with improvement in vision to 20/30, N6 (Figure 3 B).

At 1 month follow up, photoreceptoritis almost resolved with some focal disruption of ellipsoid zone in RE (Figure 2C) and LE (Figure 3C), respectively. At final visit (2 months post presentation), complete recovery with restoration of photoreceptor and outer retinal layers were seen in both eyes (Figure 2D and Figure 3D) with visual acuity improving to 20/20, N6 in both eyes with a resolution of scotoma.

DISCUSSION

This case highlights the occurrence of foveolitis following infection with dengue. A dengue foveolitis corresponds to a disruption of the outer neurosensory retina in OCT, which




Figure 3: (A) At presentation - OCT horizontal line scan through the foveal lesion shows “Conical foveal lesion”, focal outer neurosensory retina–RPE thickening. Y’ shaped split can be seen on outer plexiform layer, (B) At 1 week, comparative scan shows resolution of conical foveal lesion but photoreceptoritis still present, (C) At 1 month photoreceptoritis resolving, (D) Complete resolution seen at 2 month follow-up.

is seen clinically as yellow– orange lesions in the foveal region.⁵ Teoh *et al.* described three OCT patterns in dengue maculopathy: Type 1 (diffuse retinal thickening; 44.6%), type 2 (cystoid macular edema; 21.6%), and type 3 (foveolitis; 33.8%).⁶ Foveolitis can present with BCVA ranging from 20/20 to counting fingers. They usually present on day 7 after the onset of dengue fever.⁶ An immune-mediated mechanism may play an important role in foveolitis, similar to dengue maculopathy.^{7,8} ‘Y’ shaped split can be seen on OCT in outer plexiform layer when central macula is elevated either due to serous or solid elevation.⁹ Ophthalmologists should be aware of foveolitis associated with dengue fever which may present with acute bilateral severe visual loss. Serial OCT is useful for the diagnosis and monitoring of the disease.

REFERENCES

- Juanarita J, Azmi MN, Azhany Y, Liza-Sharmini AT. Dengue related maculopathy and foveolitis. *Asian Pacific Journal of Tropical Biomedicine*. 2012 Sep 1;2(9):755-6.
- Gupta S, Kumar A. Design of an Epitope-Based Peptide Vaccine Against Dengue Virus Isolate from Eastern Uttar Pradesh, India. *International Journal of Peptide Research and Therapeutics*. 2022 Apr 18;28(3):91.
- Yip VC, Sanjay S, Koh YT. Ophthalmic complications of dengue fever: a systematic review. *Ophthalmology and therapy*. 2012 Dec;1:1-9.
- Bacsal KE, Chee SP, Cheng CL, Flores JV. Dengue-associated

- maculopathy. *Archives of Ophthalmology*. 2007 Apr 1;125(4):501-10.
5. Loh BK, Bacsal K, Chee SP, Cheng BC, Wong D. Foveolitis associated with dengue fever: a case series. *Ophthalmologica*. 2008;222(5):317-20.
 6. Teoh SC, Chee CK, Laude A, Goh KY, Barkham T, Ang BS, Eye Institute Dengue-related Ophthalmic Complications Workgroup. Optical coherence tomography patterns as predictors of visual outcome in dengue-related maculopathy. *Retina*. 2010 Mar 1;30(3):390-8.
 7. Wen KH, Sheu MM, Chung CB, Wang HZ, Chen CW. The ocular fundus findings in dengue fever. *Gaoxiong yi xue ke xue za zhi= The Kaohsiung Journal of Medical Sciences*. 1989 Jan 1;5(1):24-30.
 8. Lim WK, Mathur R, Koh A, Yeoh R, Chee SP. Ocular manifestations of dengue fever. *Ophthalmology*. 2004 Nov 1;111(11):2057-64.
 9. Padhy SK, Kelgaonkar A, Khalsa A, Sahu S. 'Y'split of the outer plexiform layer: an optical coherence tomography illusion!. *BMJ Case Reports*. 2021 Apr 1;14(4).