Surgically Induced Focal Scleral Necrosis – A Rare Outcome

Raina Garg^{1*}, Tejal Srivastava¹, Mohammad Saquib², Abdul Waris³

¹Junior Resident, Aligarh Muslim University Institute of Ophthalmology, Aligarh, Uttar Pradesh, India.

²Assistant Professor, Aligarh Muslim University Institute of Ophthalmology, Aligarh, Uttar Pradesh, India.

³Professor, Aligarh Muslim University Institute of Ophthalmology, Aligarh, Uttar Pradesh, India.

Abstract

Purpose: To present a rare case of surgically induced scleral necrosis (SISN) following vitreoretinal surgery and discuss its clinical presentation, potential etiological factors, and management strategies.

Methods: A 71-year-old male with a history of hypertension and type 2 diabetes mellitus presented with progressive vision loss in the right eye for six months and ocular discomfort with redness for two months. Two months prior, he had undergone pars plana vitrectomy, scleral buckling, and epiretinal membrane peeling for retinal detachment. Clinical examination revealed reduced visual acuity (1/60 in OD), conjunctival congestion, scleral thinning with uveal exposure, and prominent episcleral vessels. Fundus findings included a pale, tilted optic disc with vessel shifting in the right eye, and hypertensive changes in the left eye. Autoimmune markers (ANA, RA factor) were negative. A diagnosis of post-retinal surgery-induced scleral necrosis was made.

Results: Management included topical fortified vitamin C, antibiotics, corticosteroids, and oral doxycycline. Surgical intervention with scleral patch graft and buckle readjustment was advised. The pathogenesis was attributed to possible scleral ischemia induced by excessive buckle tension or trocar-related trauma, compounded by systemic comorbidities impairing healing.

Conclusion: SISN, though rare, is a serious complication of ocular surgery. Risk factors include multiple surgeries, adjunctive therapies, surgical trauma, and systemic diseases. Early recognition and a multidisciplinary approach are critical for preserving ocular integrity. Surgeons must exercise caution during scleral manipulations and conduct thorough systemic evaluations prior to surgery.

Keywords: Scleral necrosis, Vitrectomy, Scleral buckling, Epiretinal membrane, Vitreo-retinal surgery, Non-contact tonometry, Right eye, Left eye.

INTRODUCTION

Scleral necrosis following ocular surgery is a rare but potentially devastating complication. ^{1,2} It involves localised scleral tissue death (necrosis) that can lead to severe visual morbidity. This report presents a case of Scleral necrosis induced by vitreous-retinal surgery, discusses its potential causes, clinical findings and management strategies.

Case Report

A 71-year-old male patient presented with diminution of vision in his right eye for 6 months and excessive tearing, discomfort and redness in his right eye for 2 months. He also had a history of hypertension for 20 years and type 2 diabetes mellitus for 15 years. 2 months back, he was diagnosed with retinal detachment for which he had undergone vitrectomy,

STATE OF STA

UP JOURNAL OF OPHTHALMOLOGY

An Official Journal of Uttar Pradesh State Ophthalmological Society, UPSOS (Northern Ophthalmological Society, NOS)

p-ISSN: 2319-2062 DOI: 10.56692/upjo.2025130210

scleral buckling and ERM peeling. On examination, his best corrected visual acuity (by Snellen chart) was -2.00 DS -1.00 x 110 DC 1/60 in OD and Plain 6/12 in OS. His NCT was 19 and 22 in OD and OS, respectively. Projection of rays was accurate in all quadrants, ocular movements were full in all gazes and pupillary reaction, both direct and consensual, were present in both eyes.

Address for correspondence: Raina Garg,

Junior Resident, Aligarh Muslim University Institute of Ophthalmology, Aligarh, Uttar Pradesh, India.

E-mail: rainagarg09@gmail.com

© UPJ0, 2025 Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate creative to the original author(s) and the source, provide a link to the Creative Commons licence, and incidate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit https://creativecommons.org/licenses/by-nc-sa/4.07.

How to cite this article: Garg R, Srivastava T, Saquib M, Waris A. Surgically Induced Focal Scleral Necrosis – A Rare Outcome. UP Journal of Ophthalmology. 2025;13(2): 80-82.

Received: 22-07-25, Accepted: 10-08-25, Published: 26-08-25



Figure. 1: Slit lamp picture of right eye showing retracted and congested conjunctiva, scleral thinning with exposure of the underlying uvea and prominent episcleral vessels

On slit lamp examination of the right eye, the conjunctiva was retracted and congested, scleral thinning was noted with exposure of the underlying uvea and prominent episcleral vessels (Figure 1).

Slit lamp examination of the left eye was normal. On Fundus examination of the right eye, a red glow was present, the media was clear, disc was vertically oval, tilted and pale; arterio-venous nicking was observed in the infero-temporal quadrant along with nasal shifting of blood vessels (Figures 2 and 3).

Left eye fundus examination shows soft exudates in the background, mainly near the optic disc in the supero-nasal quadrant and copper wiring of the blood vessels in the inferotemporal quadrant.

Antinuclear antibody and rheumatoid arthritis factor were found to be negative.

A diagnosis of post-retinal surgery induced scleral necrosis was made.



Figure. 2: Fundus picture of the right eye



Figure. 3: Fundus picture of left eye

Management

Patient was started on topical fortified vitamin C 4 times a day, topical moxifloxacin thrice daily, topical prednisolone 6 times a day and tablet doxycycline 1 tablet, twice daily. He was also advised on scleral patch grafting with buckling readjustment.

DISCUSSION

Surgically induced scleral necrosis is a rare but serious complication following ocular surgeries, particularly those involving the sclera or conjunctiva. 1.2 The use of adjunctive agents like mitomycin C, while effective at reducing scarring and recurrence of conditions like pterygopalatine, carries the risk of delayed wound healing and tissue necrosis. 3,4

Cryotherapy/Diathermy, often used to treat retinal tears, but overuse or high-intensity application can cause scleral damage. 5,6

In this case, excessive tension due to scleral buckling or the use of trocars might have led to scleral ischemia.⁷ Advanced age and comorbidities like diabetes or autoimmune diseases can impair wound healing.⁸

Careful placement of trocars and scleral buckles can reduce the risk of ischemia and necrosis. Early signs of scleral necrosis (pain, redness, scleral discolouration) should be identified promptly for early intervention.

Conclusion

Since surgically induced scleral necrosis is uncommon, it underscores the need for vigilance and meticulous surgical planning to minimize risks and optimize patient outcomes. Early recognition and intervention are essential to preserving vision and ocular structure.

Carry Home Message

 Undergoing multiple vitrectomies is a recognized risk factor for surgically induced scleral necrosis.

- Systemic lupus erythematosus, rheumatoid arthritis, and other collagen vascular disorders should be excluded prior to performing surgeries involving the sclera.
- A thorough systemic evaluation and gentle surgical technique are key to preventing surgically induced scleral necrosis.

REFERENCES

- Gaonker T, Yadav A, Rastogi A, Bansal P. Surgically induced necrotizing scleritis: An overview. In: Jeng BH, editor. EyeWiki [Internet]. American Academy of Ophthalmology; [cited 2025 Apr 19]. Available from: https://eyewiki.org/ Surgically-Induced Necrotizing Scleritis.
- Ruiz-Lozano RE, Garza-Garza LA, Davila-Cavazos O, Foster CS, Rodriguez-Garcia A. The clinical and pathogenic spectrum of surgically-induced scleral necrosis: A review. Surv Ophthalmol. 2021 Jul-Aug;66(4):594-611. doi: 10.1016/j. survophthal.2020.12.008. Epub 2021 Jan 8. PMID: 33422510.
- Martins TG, Costa AL, Alves MR, Chammas R, Schor P. Mitomycin C in pterygium treatment. Int J Ophthalmol. 2016 Mar 18;9(3):465-8. doi: 10.18240/ijo.2016.03.25. PMID: 27158622; PMCID: PMC4844053.
- 4. Heiligenhaus A, Akova Y, Lung E, Schrenk M, Waubke TN.

- Pterygium exzision mit intraoperativer Applikation von niedrig dosiertem Mitomycin C [Pterygium excision with intra-operative administration of low dosage mitomycin C]. Ophthalmologe. 1995 Aug;92(4):458-62. German. PMID: 7549329
- Bhatt, N., Peyman, G.A. & Karaçorlu, M. Scleral damage: comparison of standard and modified diathermy electrodes. *Int Ophthalmol* 17, 255–258 (1993). https://doi.org/10.1007/ BF01007792
- Harsum SE, Sullivan PM. Complications of scleral buckling. Retina Today. 2009 Sep;2009(9):60–61.
- Sreenivasan J, Joshi A, Chawla A. Anterior segment ischemia following scleral buckle surgery: The forgotten enemy. Oman J Ophthalmol. 2024 Oct 24;17(3):396-398. doi: 10.4103/ojo. ojo 34 24. PMID: 39651517; PMCID: PMC11620293.
- Spampinato SF, Caruso GI, De Pasquale R, Sortino MA, Merlo S. The Treatment of Impaired Wound Healing in Diabetes: Looking among Old Drugs. Pharmaceuticals (Basel). 2020 Apr 1;13(4):60. doi: 10.3390/ph13040060. PMID: 32244718; PMCID: PMC7243111.
- Jomar DE, AlHilali S, AlMutlak M. Scleral melt and uveal prolapse following 23-gauge pars plana vitrectomy. Am J Ophthalmol Case Rep. 2023;29:101769. doi:10.1016/j. ajoc.2022.101769